

Spontaneous emission and teleportation in cavity QED

P P Munhoz[†], J A Roversi, A Vidiella-Barranco and F L Semião

Instituto de Física “Gleb Wataghin” - Universidade Estadual de Campinas,
13083-970 Campinas São Paulo Brazil

Abstract. In this work, we consider atomic spontaneous emission in a system consisting of two identical two-level atoms interacting dispersively with the quantized electromagnetic field in a high-Q cavity. We investigate the destructive effect of the atomic decay on the generation of maximally entangled states, following the proposal by Zheng S B and Guo G C (2000 *Phys. Rev. Lett.* **85** 2392). In particular, we analyze the fidelity of teleportation performed using such a noisy channel and calculate the maximum spontaneous decay rate we may have in order to realize teleportation.

PACS numbers: 03.65.Yz, 03.67.Mn, 42.50.-p

1. Introduction

The generation and coherent control of maximally entangled states is of fundamental importance in the achievement of various information processing tasks as superdense coding [1] and quantum teleportation [2]. If a pure but not maximally entangled state is used to perform the standard teleportation scheme one faces two possibilities: either it is possible to teleport only particular qubit states, or teleportation is done with reduced fidelity. Some clever ideas intending to increase such a teleportation fidelity have been reported. One possibility is to concentrate partial entanglement by using local operations and classical communication [3]. In this case, the sender (Alice) and the receiver (Bob) are supplied with n pairs of identical partial entangled states and, for large n , the fraction of maximally entangled states obtained is nE , being E the von Neumann entropy of the density matrices obtained by tracing out one of the subsystems. Another interesting possibility is the development of optimal strategies that slightly change the standard protocol to maximize the probability of perfect teleportation [4]. In a more realistic situation, however, the state shared by Alice and Bob will happen to be mixed rather than pure due to unwanted noise and imperfections. In this case, it is still possible to distill out some quantum entanglement from those states. If the fidelity of the mixed state relative to a perfect singlet is larger than $\frac{1}{2}$ and Alice and Bob are

[†] Corresponding author: pmunhoz@ifi.unicamp.br

provided with many identical states, there is still a purification scheme that yields some small fraction of almost perfectly pure singlet states that can be used to high fidelity quantum teleportation [5].

The practical implementation of teleportation requires a high degree of control upon physical systems. The first experimental teleportation reported made use of correlated photon pairs produced by parametric down-conversion [6]. Since then, it has been experimentally carried out in a wide variety of quantum systems [7]. In cavity quantum electrodynamics (QED), one of the most interesting proposals is based on the dispersive interaction of two identical atoms with a single-mode cavity field (cavity-assisted atomic collisions) [8]. In the ideal case, this process would lead to the generation of perfectly entangled states (noiseless channel). From the experimental side, although teleportation using this system has not yet been performed, the coherent coupling between the atoms has already been demonstrated [9]. Experimental imperfections (three-body collisions, for instance) cause the fidelity and entanglement of the generated state to decrease and this destructive effect can indirectly be seen in experimentally accessible joint detection probabilities. In the experiment in reference [9], two Rydberg atoms cross a nonresonant cavity and become entangled by virtual emission and absorption of microwave photons. As a consequence, the procedure is supposed to be insensitive to thermal photons and cavity decay. However, one may also think of possible decoherence channels for the atoms, such as atomic spontaneous emission, as having destructive effects. If one of the atoms decays, the global atomic state factorizes and all the entanglement is lost. In this paper, we include spontaneous emission in the two-atom system interacting through a cavity field and analyze the problem of generation of maximally entangled states as well as its use in teleportation. We compute the joint detection probabilities and analyze their dependence on the atomic decaying rate for different atom-field detunings. Once the spontaneous emission leads the two-atom system to a mixed state, we find one of the main results of our paper: an upper limit for the value of the spontaneous emission rate below which one can perform quantum teleportation using that particular noisy channel. We would like to point out that although dissipation has normally destructive effects on quantum coherence, it may also allow the generation of entangled atomic states [10] and nonclassical states of the vibrational motion of a trapped ion [11]. In other studies it is shown that an adequate driving of the system may compensate losses, yielding the stabilization of entanglement between two atoms [12].

2. Solution of the Master Equation

We consider a system consisting of two identical two-level atoms interacting with a quantized electromagnetic field enclosed in a high- Q cavity [13], as shown in figure (1). If the two atoms are sufficiently far from each other, we can neglect the dipole-dipole interaction and the Hamiltonian for this system in the interaction picture reads [8, 14]

$$\mathbf{H} = \hbar\lambda \sum_{j=1,2} (e^{-i\delta t} \mathbf{a}^\dagger \boldsymbol{\sigma}_j + e^{i\delta t} \mathbf{a} \boldsymbol{\sigma}_j^\dagger), \quad (1)$$




Figure 1. A sketch of the system comprising a couple of two-level atoms (frequencies ω_a), in a far detuned high-Q cavity (frequency ω_f). The atoms interact with each other via the effective cavity-assisted atom-atom coupling Ω . We assume that the decay rate γ is the same for each atom.

where λ is the atom-cavity coupling constant, σ_j ($\sigma_j^\dagger$) is the atomic lowering (raising) operator, $\mathbf{a}$ ($\mathbf{a}^\dagger$) is the annihilation (creation) operator of photons in the field mode and δ the detuning between the atomic and cavity field frequencies denoted by ω_a and ω_f , respectively. As considered in the original maximally entangled state generation proposal [8], we take here the dispersive limit ($\delta \gg \lambda$) in Hamiltonian (1). If there are no photons initially in the cavity (vacuum state), the system may be described by an effective two-atom Hamiltonian which, in the interaction picture, is given by [8]

$$H = \hbar\Omega(\sigma_1^\dagger\sigma_2 + \sigma_2^\dagger\sigma_1 + \sigma_1^\dagger\sigma_2 + \sigma_1\sigma_2^\dagger), \quad (2)$$

where $\Omega = \lambda^2/\delta$ is the effective cavity-assisted atom-atom coupling constant.

The above Hamiltonian governs the dynamics of the system in the ideal case where losses are not included. The Hamiltonian (2) might lead to the generation of a maximally entangled state. Consider, for instance, that atom 1 is initially prepared in the state $|e_1\rangle$ and the atom 2 in the state $|g_2\rangle$, i.e., the initial global state of the two atoms is $|\psi(0)\rangle = |e_1\rangle \otimes |g_2\rangle \equiv |eg\rangle$. In this case, the evolved state will be [8]

$$|\psi(t)\rangle = e^{-i\Omega t}[\cos(\Omega t)|eg\rangle - i\sin(\Omega t)|ge\rangle]. \quad (3)$$

For interaction times[‡] $t_k = (2k+1)\pi/4\Omega$ (k integer) that corresponds to the maximally entangled state

$$|\psi_{\text{EPR}}\rangle = \frac{1}{\sqrt{2}}(|eg\rangle - i|ge\rangle), \quad (4)$$

which can be used for faithful teleportation as reported in [8].

[‡] Or equivalently, detunings $\delta_k = 4\lambda^2 t/(2k+1)\pi$.

However, in a more realistic situation, mechanisms of losses may be present. Two of the most important dissipation channels are the leaking of photons through the cavity walls and the atomic spontaneous emission of photons into noncavity field modes. In general, losses can be described as the irreversible coupling of the system to a large reservoir. In the zero temperature case, and considering the cavity field initially in the vacuum state, the losses due to cavity damping are irrelevant according to the effective Hamiltonian (2). For this reason, we consider here just the atomic spontaneous emission which is expected to degrade the maximum amount of entanglement in state (4) until the system reaches its ground state $|gg\rangle$ (disentangled state). In the rotating wave and Born-Markov approximations, the density operator for two atoms obeys the following master equation [15,16]

$$\dot{\rho}(t) = \frac{1}{i\hbar}[\mathbf{H}, \rho(t)] + \mathcal{L}_a \rho(t), \quad (5)$$

where

$$\mathcal{L}_a = 2\gamma(\sigma_1 \cdot \sigma_1^\dagger + \sigma_2 \cdot \sigma_2^\dagger - \frac{1}{2}\{\sigma_1^\dagger \sigma_1 + \sigma_2^\dagger \sigma_2, \cdot\}), \quad (6)$$

and γ is the spontaneous emission rate[§]. We have used the superoperator notation $\mathbf{O} \cdot (\cdot \mathbf{O})$ which represents the action of an operator $\mathbf{O}$ to the left (right) on the target operator and this convention will be used hereafter. Once we are considering that the atoms are not close enough to take their mutual interaction into account, the atom-atom cooperation induced by their coupling with a common reservoir is not included in master equation (5).

As we show next, one can exactly solve (5) by means of the application of the following unitary transformation [17]

$$\mathbf{U} = \exp \left[-\frac{\pi}{4}(\sigma_1^\dagger \sigma_2 - \sigma_1 \sigma_2^\dagger) \right]. \quad (7)$$

The operator $\mathbf{U}$ commutes with $\mathbf{N} = \sigma_1^z + \sigma_2^z$, i.e., $[\mathbf{U}, \mathbf{N}] = 0$, i.e., the transformation above preserves the total excitation number $\mathbf{N}$, simplifying the solution of the problem in cases where the initial excitation number is well defined. In this work we will consider the case $N = 0, 1$ because it is consistent with the initial preparation considered in [8]. Now we may restrict ourselves to the Hilbert subspace spanned by the basis $\{|g_1 g_2\rangle, |e_1 g_2\rangle, |g_1 e_2\rangle\}$, and move to a frame according to

$$\mathbf{V} = \exp \left[i\Omega(\sigma_1^\dagger \sigma_1 + \frac{1}{2}\sigma_1^z)t \right]. \quad (8)$$

We can rewrite the master equation (5) as

$$\dot{\tilde{\rho}}(t) = \mathbf{T}^\dagger \dot{\rho}(t) \mathbf{T} = \mathcal{L}_a \tilde{\rho}(t), \quad (9)$$

where $\mathbf{T} = \mathbf{U}\mathbf{V}$ and, after defining the superoperators

$$\begin{aligned} \mathbf{J} &= \mathbf{J}_1 + \mathbf{J}_2, & \mathbf{J}_i &= \sigma_i \cdot \sigma_i^\dagger, \\ \mathbf{L} &= \mathbf{L}_1 + \mathbf{L}_2, & \mathbf{L}_i &= -\frac{1}{2}\{\sigma_i^\dagger \sigma_i, \cdot\}, \end{aligned} \quad (i = 1, 2) \quad (10)$$

[§] For simplicity we assume that both atoms are of the same type and are placed inside the cavity in positions in which the electric field is basically the same. Therefore the atoms should have equal spontaneous emission rates.

the master equation (9) assumes the simple form

$$\dot{\tilde{\rho}}(t) = 2\gamma(\mathbf{J} + \mathbf{L})\tilde{\rho}(t). \quad (11)$$

Its solution may be written as [18]

$$\tilde{\rho}(t) = \exp[(e^{2\gamma t} - 1)\mathbf{J}] \exp(2\gamma t\mathbf{L})\tilde{\rho}(0), \quad (12)$$

where $\tilde{\rho}(0) = \mathbf{T}^\dagger \rho(0) \mathbf{T}$. Considering the initial state to be $\rho(0) = |eg\rangle\langle eg| = \sigma_1^\dagger \sigma_1 \sigma_2 \sigma_2^\dagger$, we have

$$\tilde{\rho}(0) = \frac{1}{2}(\sigma_1^\dagger \sigma_1 \sigma_2 \sigma_2^\dagger - \sigma_1^\dagger \sigma_2 - \sigma_1 \sigma_2^\dagger + \sigma_1 \sigma_1^\dagger \sigma_2^\dagger \sigma_2), \quad (13)$$

and then, the evolved state according to (11) is given by

$$\tilde{\rho}(t) = \frac{1}{2}e^{-2\gamma t}[\sigma_1^\dagger \sigma_1 \sigma_2 \sigma_2^\dagger - \sigma_1^\dagger \sigma_2 - \sigma_1 \sigma_2^\dagger + \sigma_1 \sigma_1^\dagger \sigma_2^\dagger \sigma_2 + 2(e^{2\gamma t} - 1)\sigma_1 \sigma_1^\dagger \sigma_2 \sigma_2^\dagger]. \quad (14)$$

It is not difficult to show that, when transformed back to the original space, the state (14) reduces to (3) in the ideal case ($\gamma = 0$). From the density operator $\tilde{\rho}$, all the statistical properties of the system may be readily obtained||.

3. Dynamics in the Lossy Case

Now, we turn to the study of the effect of atomic spontaneous emission on the system dynamics. We focus on two quantities that may be readily measured in this system [9, 19].

3.1. Joint Detection Probabilities

The first quantity is the probability of detecting the two atoms in one of the four states $\{|gg\rangle, |ge\rangle, |eg\rangle, |ee\rangle\}$. Once there is no pumping mechanism acting upon the system, the state $|ee\rangle$ remains unpopulated. The joint probabilities for the initial preparation (13) can be computed from (14) and they are given by

$$P(e_1, g_2) = e^{-2\gamma t} \cos^2(\Omega t), \quad (15)$$

$$P(g_1, e_2) = e^{-2\gamma t} \sin^2(\Omega t), \quad (16)$$

$$P(g_1, g_2) = 1 - e^{-2\gamma t}. \quad (17)$$

In figure(2), we show these probabilities as a function of the atom-field detuning following the lines in [9], where the experimental curve is presented. In figure 2, one may see that the spontaneous emission plays an important role in the joint detection probabilities. In the ideal case ($\gamma/\lambda = 0$), i.e., the absence of spontaneous emission, it happens that for times when $P(e_1, g_2)$ is equal to $P(g_1, e_2)$, it occurs the generation of the EPR state (4) and neither $|gg\rangle$ nor $|ee\rangle$ is populated. The situation is quite different when one considers the more realistic setup where the atoms may decay spontaneously. Although $P(e_1, g_2)$ and $P(g_1, e_2)$ still cross each other, their amplitudes become now

|| One should remember that for a given observable operator $\mathbf{A}$, its expectation value $\langle \mathbf{A} \rangle$ is given by either $\text{Tr}[\rho \mathbf{A}]$ or $\text{Tr}[\tilde{\rho} \tilde{\mathbf{A}}]$, being the latter referred to any space transformed by unitary operations.




Figure 2. Joint probabilities versus the ratio λ/δ . $P(g_1, g_2)$, $P(e_1, g_2)$, and $P(g_1, e_2)$ are represented by solid, dashed, and dotted lines, respectively. For all plots, λt is set equal to 3π . (a) $\gamma/\lambda = 0$; (b) $\gamma/\lambda = 0.01$; (c) $\gamma/\lambda = 0.05$; (d) $\gamma/\lambda = 0.1$.

suppressed due to the fact that the stronger the decay rate γ , the more important is the mixture with the component $|gg\rangle$. Not surprisingly, that must be the case, because the system ground state is expected to become more populated as the atomic levels couple strongly to the reservoir. When comparing those results to the experimental curves, one immediately finds common features. As pointed out in [9], imperfections such as erroneous detection counts or a third atom crossing the cavity, both could lead to non null spurious probabilities $P(g_1, g_2)$ and $P(e_1, e_2)$ and, consequently, to the suppression of joint probabilities involving other states [9, 20]. However, according to what it is shown here, even if those experimental imperfections were fixed, e.g., by allowing just two atoms inside the cavity at a time and by improving the detection efficiency, it would still be a spurious population in the ground state due to atomic spontaneous emission. For the same reason the quantity $P(e_1, g_2)$ would be suppressed in the limit $\lambda/\delta \rightarrow 0$ (infinite detuning). Of course, these effects become less important as long lived atomic levels are chosen. It is noteworthy that for small detunings one should use the original Hamiltonian (1) rather than the effective (dispersive) one given by equation (2). For the purpose of the generation of the maximally entangled state, however, one is interested in the dispersive limit, that starts to be a good approximation for $\delta > 3\lambda$ [9].




Figure 3. Bell signal as a function of the angular variable ϕ . For all plots, Ωt is set equal to $\pi/4$. We considered the ratio $\gamma/\Omega = 0$ (solid), $\gamma/\Omega = 0.2$ (dashed), and $\gamma/\Omega = 0.4$ (dotted).

3.2. Bell Signal Analysis

The second quantity of interest is the expectation value $\beta(\phi) \equiv \langle \sigma_1^x \sigma_2^\phi \rangle$, where $\sigma_i^\phi = \cos(\phi)\sigma_i^x + \sin(\phi)\sigma_i^y$, being σ_i^x and σ_i^y Pauli matrices for the atom i . Fixing the interaction time to be $t_k = (2k + 1)\pi/4\Omega$ (k integer), the variation of the Bell signal (as a function of ϕ) measures the angular correlations between the transverse components associated to the atoms. Consequently, this quantity gives a measure of the purity of the generated state. It is easy to show that an equally weighted mixture of the states $|eg\rangle$ and $|ge\rangle$ would give $\langle \sigma_1^x \sigma_2^\phi \rangle = 0$ for all ϕ . In this sense, looking at the joint detection probabilities alone is not enough to decide if a pure EPR state has been generated. In other words, the joint probabilities give information about the diagonal elements of the density matrix whereas the Bell signal furnishes information about its non diagonal elements. This quantity can be computed directly from (14) and the result is

$$\beta(\phi) = e^{-\gamma(2k+1)\pi/2\Omega} \cos(\phi). \quad (18)$$

Again, the effect of spontaneous emission is to cause a suppression of the oscillations as shown in figure (3). The same behaviour was experimentally observed [9].

4. Fidelity of Teleportation

In the standard teleportation protocol [2], two parties Alice and Bob are supplied with a pair of particles in a EPR state. They keep their respective part with one another to function as a quantum channel to teleport a third particle in possession of Alice, which is in an unknown pure state $|\psi\rangle = a|e\rangle + b|g\rangle$. Alice then separately performs a joint measurement on her part of the EPR pair and on the state she wants to




Figure 4. Scheme of quantum teleportation of an unknown quantum state $|\psi\rangle$. Alice and Bob share a previously generated quantum channel, a EPR pair. After combining the state $|\psi\rangle$ with her portion of the EPR pair, Alice performs a Bell state measurement, destroying the original state. She then informs Bob the result of her measurement, so that he may perform the correct operation on his portion of the EPR pair and thus recover the original teleported state $|\psi\rangle$.

teleport, and sends the result of her measurement to Bob using a classical channel (e.g., a phone call). As the quantum channel is maximally entangled, no matter what outcome from the Bell measurement Alice gets, Bob will have his part of the channel projected into one of four possible transformations of the original but no longer existing pure state $|\psi\rangle$. To accomplish the teleportation procedure, Bob only needs to apply, based on the information received from Alice, a combination of the four transformations ($\mathbf{I}, \sigma^x, \sigma^y, \sigma^z$). The protocol is depicted in figure (4).

Maximally entangled states provide a quantum channel that allows the perfect teleportation of an unknown state from one party to another. After the seminal paper [2], it seemed natural to search for connections between teleportation, Bell's inequalities and inseparability [21–24]. One of the first things that had been noticed was the fact that a mixed state could still be useful for (imperfect) teleportation [21]; it was demonstrated that the fidelity of transmission of an unknown qubit can be linked to the nonclassical character of the state forming the quantum channel. Actually, a classical channel can give at most a fidelity equal to $\frac{2}{3}$. This may be achieved if Alice simply performs a measurement on the unknown state and tells the result to Bob [21].

In the system treated here, the two atoms become maximally entangled at specific interaction times when no imperfections or losses are considered in the model. Then, a practical question that naturally arises is: What is the maximum value of the decay constant γ that still allows legitimate, although imperfect quantum teleportation? We answer that question using the fidelity of teleportation discussed in [22]. They showed that the maximum fidelity that can be obtained using a noisy channel ρ is given by

$$F_{\max} = \frac{1}{2} \left(1 + \frac{1}{3} \text{Tr} \sqrt{T^\dagger T} \right), \quad (19)$$




Figure 5. Fidelity of teleportation as a function of the scaled time Ωt and decay rate γ/Ω .

where T is a real matrix formed by the elements $t_{nm} = \text{Tr}[\rho(\sigma_1^n \otimes \sigma_2^m)]$, where $\{\sigma_i^n\}_{n=1}^3$ are the ordered Pauli matrices $\{\sigma_i^x, \sigma_i^y, \sigma_i^z\}$ for the atom i . In the interaction times t_k , the maximum fidelity of teleportation may be obtained from (14) and it is given by

$$F_{\max} = \begin{cases} \frac{1}{3} + \frac{2}{3}e^{-\gamma(2k+1)\pi/2\Omega}, & \text{for } \gamma/\Omega \leq \ln(4)/(2k+1)\pi \\ \frac{2}{3}, & \text{otherwise.} \end{cases} \quad (20)$$

For those interaction times, a fidelity of $\frac{2}{3}$ indicates that the system has decayed to a state that no longer allows reliable quantum teleportation. The higher the γ , the closer the system state gets closer to the ground state $|gg\rangle$. Now, we can answer the question raised above: the system provides a quantum channel suitable for teleportation, at interaction times t_k , provided the decay constant γ does not exceed the upper limit $\gamma = \Omega \ln(4)/(2k+1)\pi$.

Of course, it is not just for the specific interaction times t_k that one would obtain $F_{\max} > \frac{2}{3}$. Precisely at those times the function $F_{\max}$ has local maxima. This can be seen in figure (5), where there is a plot of $F_{\max}$ as a function of the scaled time and the decay rate.

5. Entanglement

In this section we analyze the amount of entanglement between the two atoms using the concurrence, defined as [25]

$$C(\rho) = \max\{0, \lambda_1 - \lambda_2 - \lambda_3 - \lambda_4\}, \quad (21)$$

where λ_i are the square roots of the eigenvalues, in decreasing order, of the non-Hermitian matrix $\rho\rho_{\text{sf}}$, being

$$\rho_{\text{sf}} = (\sigma_1^y \otimes \sigma_2^y)\rho^*(\sigma_1^y \otimes \sigma_2^y), \quad (22)$$

the “spin flip” transformation of the complex conjugation of ρ in the standard basis $\{|ee\rangle, |eg\rangle, |ge\rangle, |gg\rangle\}$, and σ_i^y is the Pauli spin matrix. For a maximally entangled state $C(\rho) = 1$, while for a separable state $C(\rho) = 0$.




Figure 6. Concurrence as a function of the scaled time Ωt and decay rate γ/Ω .

At the interaction times t_k , the concurrence is given by

$$C(\rho) = e^{-\gamma(2k+1)\pi/2\Omega}, \quad (23)$$

which is related to the maximum fidelity of teleportation as

$$F_{\max} = \begin{cases} \frac{1}{3} + \frac{2}{3}C(\rho), & \text{for } \gamma/\Omega \leq \ln(4)/(2k+1)\pi \\ \frac{2}{3}, & \text{otherwise.} \end{cases} \quad (24)$$

Therefore, in this case, for $C(\rho) \geq \frac{1}{2}$ the system provides a quantum channel for reliable teleportation. The dependence of entanglement on time and on the damping rate is shown in figure (6), where we have a plot of the concurrence as a function of Ωt and the decay rate γ/Ω . In comparing figure (6) with figure (5) one may find, for instance, ranges of parameters for which quantum teleportation is not allowed despite the existence of entanglement.

6. Conclusions

We have investigated the effect of atomic spontaneous emission in the generation of maximally entangled states, which are useful for quantum teleportation. We have shown that experimental observables such as joint probabilities and correlation functions involving transverse components of the Pauli matrices both are suppressed due to the effect of atomic decay. We have computed the fidelity of teleportation and found an upper limit for the value of the spontaneous emission rate below one can perform quantum teleportation using that noisy channel. The knowledge of the upper bound for the decay rate allowing the realization of teleportation, as we have found here, is of central importance if one wants to implement such a quantum channel in an actual experiment. The effect considered here may be lessened if the spontaneous emission is set to be negligible, what would correspond to the choice of long lived atomic levels. Nevertheless, if one wants to address a more general situation, the effects of spontaneous emission are not negligible and have to be taken into account. We have also calculated the amount of entanglement (quantified by the concurrence) between the

atoms as a function of the atomic decay rate, which shows that a substantial amount of entanglement is required in order to allow quantum teleportation using the atomic quantum channel here discussed.

Acknowledgments

This work is partially supported by CNPq (Conselho Nacional para o Desenvolvimento Científico e Tecnológico), and FAPESP (Fundação de Amparo à Pesquisa do Estado de São Paulo) grant number 02/02715-2, Brazil.

References

- [1] Bennett C H and Wiesner S J 1992 *Phys. Rev. Lett.* **69** 2881
- [2] Bennett C H *et al* 1993 *Phys. Rev. Lett.* **70** 1895
- [3] Bennett C H *et al* 1996 *Phys. Rev. A* **53** 2043
- [4] Li W L *et al* 1996 *Phys. Rev. A* **61** 034301
- [5] Bennett C H *et al* 1996 *Phys. Rev. Lett.* **76** 722
- [6] Bouwmeester B *et al* 1997 *Nature* **390** 575
- [7] Boschi E *et al* 1998 *Phys. Rev. Lett.* **80** 1121
 - Furusawa A *et al* 1998 *Science* **282** 706
 - Nielsen M A *et al* 1998 *Nature* **396** 52
 - Pan J W *et al* 2001 *Phys. Rev. Lett.* **86** 4435
 - Marcikic I *et al* 2002 *Nature* **421** 509
 - Fattal D *et al* 2004 *Phys. Rev. Lett.* **92** 037904
 - Riebe M *et al* 2004 *Nature* **429** 734
 - Barret M D *et al* 2004 *Nature* **429** 737
- [8] Zheng S B and Guo G C 2000 *Phys. Rev. Lett.* **85** 2392
- [9] Osnaghi S *et al* 2001 *Phys. Rev. Lett.* **87** 037902
- [10] Plenio M B *et al* 1999 *Phys. Rev. A* **59** 2468
- [11] Semião F L and Vidiella-Barranco A. 2005 *Phys. Rev. A* **71** 065802
- [12] Çakir Ö *et al* 2005 *Phys. Rev. A* **71** 034303
- [13] Ye L and Guo G C 2005 *Phys. Rev. A* **71** 034304
 - Xu J B and Li S B 2005 *New J. Phys.* **7** 72
- [14] Tavis M and Cummings F W 1968 *Phys. Rev.* **170** 379
 - Dicke R 1954 *Phys. Rev.* **93** 99
- [15] Walls D F and Milburn G J 1994 *Quantum Optics* (Berlin: Springer)
- [16] Gardiner C W and Zoller P 2000 *Quantum Noise* (Berlin: Springer)
- [17] Nicolosi S *et al* 2004 *Phys. Rev. A* **70** 022511
- [18] Barnett S M and Radmore P M 1997 *Methods in Theoretical Quantum Optics* (Oxford: Clarendon)
- [19] Brune M *et al* 1996 *Phys. Rev. Lett.* **77** 4887
- [20] Guo G P *et al* 2002 *Phys. Rev. A* **65** 042102
- [21] Popescu S 1994 *Phys. Rev. Lett.* **72** 797
- [22] Horodecki R *et al* 1996 *Phys. Lett. A* **222** 21
- [23] Massar S and Popescu S 1995 *Phys. Rev. Lett.* **74** 1259
- [24] Gisin N 1996 *Phys. Lett. A* **210** 157
 - Gisin N 1996 *Phys. Lett. A* **210** 151
- [25] Wootters W K 1998 *Phys. Rev. Lett.* **80** 2245